

Integrated Pest Management

Know 10 Common Pests of Edible Plants

About Community In Bloom

Community In Bloom (CIB) is a programme that was launched by the National Parks Board (NParks) in 2005. It aims to nurture a gardening culture among Singaporeans by encouraging and facilitating community gardening efforts. It is also an opportunity to build community bonds and strengthen social resilience.



   nparksbuzz
Share your love for nature with us at
#nparksbuzz

For more information, visit our website at
www.nparks.gov.sg/cib or email us at
CommunityInBloom@nparks.gov.sg

For more information on plants in Singapore,
visit NParks Flora & Fauna Web at
florafaunaweb.nparks.gov.sg

Methods that can be integrated into a Pest Management programme:

- 
 - Setting up sticky traps to trap flying insect pests
 - Setting up pheromone traps
 - Wrapping fruits with paper or plastic to prevent insects from laying eggs
 - Use of nets or netted structures over planting beds or plants to physically prevent the pests from coming into contact with the plants
 - Removal of larger pests such as beetles, caterpillars and snails by hand

This includes:

fences and nets to prevent pest activity or to remove pests from an area

Mechanical Control

- Neem oil and other horticultural oils
 - DIY non-chemical pesticides such as chili, garlic and onion spray
 - Attracting natural predators such as ladybugs and lacewings

This includes:

The use of natural chemicals and existing predators in the garden to control pest populations

- Altering the environment, the condition of the host plant, or the behavior of the pest to prevent or suppress an infestation or infection
- Checking for and removing pests regularly
- Growing plants in clean soil
- Providing optimal amounts of water and sunlight for plants
- Applying fertilisers appropriately
- Ensuring good air circulation around plants (frequent pruning to thin out dense shrubs)
- Removing infected plant parts promptly
- Bags and disposing of them away and properly by placing them in
- From the garden

This includes:

- Checking for and removing pests regularly
 - Growing plants in clean soil
 - Providing optimal amounts of water and sunlight for plants
 - Applying fertilisers appropriately
 - Ensuring good air circulation around plants (requent pruning to thin out dense shrubs)
 - Removing infected plant parts promptly and properly by placing them in bags and disposing of them away from the garden

Luis Junceda

Altering the environment, the condition of the host plant, or the behavior of the pest to prevent or suppress an infestation or infection


Methods that can be integrated into an Integrated Pest Management program.

- Environmentally friendly
- Unlike most pesticides animals, unlike most pesticides does not affect non-targeted
- Does not affect non-targeted
- Less harmful to the ecosystem
- Mandegemem programme;

Benefits of an Integrated Pest Management Program

is a strategy that focuses on the long-term prevention of pests through a combination of techniques such as biological control, change of cultural practices and/or mechanical control. Whenever possible, these techniques should be used on pests when they are at their most susceptible, to maximise effectiveness.

Diamondback Flea Beetles



Commonly known as Awl Snails, Allopeas snails have a yellow body and an elongated shell.

Awl snails are usually found in large numbers. They are commonly found in wet, shaded areas of greenhouses, in leaf debris, under stones and in the soil beneath grasses.

Vulnerable Plants

Most leafy vegetables and fruiting vegetables such as zucchini

Symptoms

Awl snails eat leaves, shoots and flowers, creating holes.

Preventive and Control Measures

Requires an integrated management approach, including the selection and planting of non-infected nursery stock, and manually removing the snails.

Leaf miner larvae feed on leaves, creating tunnelling mine trails. The most common leaf miners present in Singapore are *Liriomyza sativae* and *L. trifolii*.

Vulnerable Plants

Most leafy vegetables and fruiting vegetables, including beans, lady's finger and sweet pea

Symptoms

Infested leaves have translucent leaf mining trails. Infected leaves should be physically removed.

Preventive and Control Measures


Remove infested plants from the garden. As the insects may have laid eggs in the soil, it is advisable to replace the top 10 to 15 cm of existing soil as well.

Preventive and Control Measures

Trap crops, such as mustard and radish can be planted as buffer plants to attract the flea beetles away from the main leafy vegetables.

Rotating crops with leafy vegetables such as bayam and kangkong may help break the life cycle of flea beetles.

Awl Snails



The caterpillars of the Diamondback Moth are common and widespread in hot and dry weather. They are light brown to green. When disturbed, they wriggle rapidly and drop from the leaves on silk threads which they use to climb back.

Vulnerable Plants

Most leafy Brassicas, such as bai cai, caixin, kailan, watercress and xiao bai cai

Symptoms

The affected leaves can become skeletonised by the caterpillars, which are voracious feeders in their last stage.

Preventive and Control Measures

Nets can be used to cover susceptible plants. Rotating crops with non-Brassica plants, such as bayam, kangkong or onions, may help to break the insect life cycle. Infested leaves may harbour larvae or eggs. Discard the waste and do not use it for compost to prevent re-introduction to new plants.

Diamondback Flea Beetles



Adult flea beetles are small, nocturnal insects that are voracious feeders of young leafy vegetables.

Vulnerable Plants

Mostly leafy vegetables from the genus *Brassica*

Symptoms

Round holes can be found on infected leaves, and plants may display reduced growth.

Preventive and Control Measures

Remove infested plants from the garden. As the insects may have laid eggs in the soil, it is advisable to replace the top 10 to 15 cm of existing soil as well.

Preventive and Control Measures

Trap crops, such as mustard and radish can be planted as buffer plants to attract the flea beetles away from the main leafy vegetables.

Rotating crops with leafy vegetables such as bayam and kangkong may help break the life cycle of flea beetles.

Root-Knot Nematodes



Root-knot nematodes are the most widespread and serious parasitic nematode pest of plants in tropical and subtropical regions. They are infective in their larval stage

Vulnerable Plants

Leafy vegetables such as bayam and kangkong, and fruiting vegetables such as brinjal, chilli, cucumber, lady's finger, melons, radish, sweet potato and tomato

Symptoms

Severe infection can cause the roots to be badly galled, hindering water and nutrient uptake. Root decay and leaf wilting can be observed, and eventually the plant may die.

Preventive and Control Measures

No single management practice is available to control or eradicate this nematode, but crop rotation with resistant varieties or non-host plants may help to prevent infection.

Red spider Mites



Red spider mites are found primarily on the underside of infected leaves. They feed by penetrating the plant tissue with their mouthparts.

Vulnerable Plants

Leafy vegetables such as bayam and kangkong, and fruiting vegetables such as brinjal, chilli, cucumber, lady's finger, melons, radish, sweet potato and tomato

Symptoms

The leaves may appear bronze or yellow, and may curl. The fruits may show corky tissue.

Preventive and Control Measures

Use of organic soap or mineral oil can reduce infestation of red spider mites.

Introducing predators such as ladybirds and lacewing larvae can help to regulate spider mite populations.

Whiteflies



Whiteflies are often found on the undersides of the leaves. They excrete honey dew, which attracts both ants and sooty mould fungi.

Vulnerable Plants

Leafy vegetables such as bayam, and fruiting vegetables such as brinjal, chilli, lady's finger, melons, radish, sweet potato and tomato

Symptoms

Infested plants may show a variety of symptoms, including leaf or vein yellowing, yellow mosaic patterns on the leaves, and leaf curling. Sooty mould may also be present.

Preventive and Control Measures

Removal of heavily infested plants will help to prevent migration of whiteflies onto nearby plants.

Remove weeds regularly to reduce the whitefly population. Yellow sticky traps may be used to trap adult whiteflies.

Mealy Bugs



Mealy bugs can be found on any part of a plant, including the stems, flowers and the underside of leaves. They prefer warm and moist conditions.

Vulnerable Plants

Most leafy vegetables and fruiting vegetables

Symptoms

A cottony covering may be observed on affected parts of the plant.

Preventive and Control Measures

Affected plant parts can be removed, and then the plant should be closely monitored to ensure that it is free of mealy bugs.

Ants protect mealy bugs from natural enemies such as ladybirds, so controlling ant populations can help to control mealy bug infestations.