

## ***Phlogacanthus magnus*, a new combination in the Acanthaceae from Malaysia**

Y.F. Deng

Key Laboratory of Plant Resources Conservation and Sustainable Utilization,  
South China Botanical Garden, Chinese Academy of Sciences,  
Guangzhou 510650, China  
yfdeng@scbg.ac.cn

**ABSTRACT.** An examination of the type material of *Gymnostachyum magnum* C.B. Clarke (Acanthaceae) from Malaysia reveals that it is a *Phlogacanthus* rather than a *Gymnostachyum* due to having two staminodes in addition to the two fertile stamens and muticous anther-thecae. The new combination, *Phlogacanthus magnus* (C.B. Clarke) Y.F. Deng is therefore proposed.

**Keywords.** Acanthaceae, *Gymnostachyum*, Malaysia, *Phlogacanthus*

### **Introduction**

*Gymnostachyum* Nees is a genus of about 30 species of Acanthaceae distributed in the Asian tropics (Mabberley, 2008; Hu et al., 2011). It belongs to tribe Andrographideae together with *Andrographis* Nees, *Phlogacanthus* Nees, *Cystacanthus* T. Anderson, *Diotacanthus* Benth., *Graphandra* Imlay, *Haplanthodes* Kuntze and *Indoneesiella* Sreem. (Lindau, 1895; Scotland, 1992; Scotland & Vollesen, 2000; Stevens, 2014; McDade et al., 2008).

For the Malay Peninsula, Clarke (1908) recognised eleven species of *Gymnostachyum*, including ten new species, and Ridley (1923) recognised 13 species. They also suggested that *Phlogacanthus* and *Gymnostachyum* might be united into a single genus but this is in part because they confused the circumscription of *Gymnostachyum* and *Phlogacanthus*. *Phlogacanthus*, a genus of about 35 species distributed in tropical regions of Asia, can be easily distinguished from *Gymnostachyum* by having two staminodes, in addition to the two fertile stamens, and muticous anther-thecae (Hu et al., 2011; Xia & Deng, 2013). In *Gymnostachyum*, the staminodes are absent and one or both of anther-thecae are mucronate at the base. The separation of *Phlogacanthus* from *Gymnostachyum* is also supported by molecular evidence (McDade et al., 2008). During a visit to Singapore Botanic Gardens' Herbarium in 2013, I had the opportunity to examine the type material of *Gymnostachyum magnum* C.B. Clarke, one of species described by Clarke (1908). *Gymnostachyum magnum* has two stamens, two staminodes and muticous anther-thecae, characters that conform well to *Phlogacanthus*. Therefore, the new combination, *Phlogacanthus magnus* (C.B. Clarke) Y.F. Deng, is proposed below.

***Phlogacanthus magnus*** (C.B.Clarke) Y.F.Deng, **comb. nov.** – *Gymnostachyum magnum* C.B.Clarke, *J. Asiat. Soc. Bengal*, Pt. 2, Nat. Hist. 74(3): 664 (1908). TYPE: Malaysia, Negri Sembilan, Bukit Tampin waterfall, May 1894, *J.S. Goodenough 1893* (holotype K; isotype SING!).

*Distribution and habitat.* This species is endemic to Peninsular Malaysia. Its habitat details are not well known.

*Provisional IUCN Conservation Assessment.* Currently, this species is only known from three localities in Peninsular Malaysia comprising only five, mostly fairly old, collections, but no attempt has yet been made to assess its frequency in the field. Given the lack of knowledge of the current status of the species it should be assessed as Data Deficient (DD) (IUCN, 2001, 2011).

*Additional specimens examined:* PENINSULAR MALAYSIA. **Kedah:** Katanh, Koh Mai Forest Reserve, 4 Apr 1938, *Kiah* s.n. (SING). **Negeri Sembilan:** Sungei Ujong, Bukit Sutu, 1 Nov 1885, *V.M. Alvins 1955* (SING). **Selangor:** Kajang, Sungei Lalauy, 9 Mar 1930, *C.F. Symington 22710* (K); *ibidem*, without date, *C.F. Symington 22763* (SING); *ibidem*, 28 Mar 1931, *C.F. Symington 22947* (SING).

ACKNOWLEDGEMENTS. The author would like to thank Dr Wong Khoon Meng, Mr Low Yee Wen and Ms Serena Lee for their help during my visit to the Herbarium of Singapore Botanic Gardens (SING). This work was supported by the National Natural Science Foundation of China (grant nos. 31070175, 31270247, 31470302) and the Singapore Botanic Gardens' Fellowship.

## References

- Clarke, C.B. (1908). Acanthaceae. In: King, G. & Gamble, J.S. Materials for a Flora of the Malayan Peninsula. *J. Asiat. Soc. Bengal*, Pt. 2, Nat. Hist. 74: 628–698.
- Hu, J.Q., Deng, Y.F. & Wood, J.R.I. (2011). Acanthaceae. In: Wu, C.Y., Raven, P. & Hong, D.Y. (eds) *Flora of China* 19: 369–477. Beijing: Science Press & St. Louis: Missouri Botanical Garden Press.
- IUCN (2001). *IUCN Red List Categories and Criteria, Version 3.1*. Switzerland, Gland and UK, Cambridge: IUCN.
- IUCN (2011). *Guidelines for using the IUCN Red List Categories and Criteria, Version 9*. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed on 29 Jun. 2014).
- Lindau, G. (1895). Acanthaceae. In: Engler, A. (ed), *Die Natürliche Pflanzenfamilien* 4(3b): 274–354. Leipzig.
- Mabberley, D.J. (2008). *Mabberley's Plant-book: a Portable Dictionary of Plants, their Classifications, and Uses*. Cambridge: Cambridge University Press.
- McDade, L.A., Daniel, T.F. & Kiel, C.A. (2008). Toward a comprehensive understanding of phylogenetic relationships among lineages of Acanthaceae s.l. (Lamiales). *Amer. J. Bot.* 95: 1136–1152.

- Ridley, H.N. (1923). Acanthaceae. In: Ridley, H.N. (ed) *The flora of the Malay Peninsula 2*: 554–610. London: L. Reeve & Co.
- Scotland, R.W. (1992). Pollen morphology of Andrographideae (Acanthaceae). *Rev. Palaeobot. Palynol.* 72: 229–243.
- Scotland, R.W. & Vollesen, K. (2000). Classification of Acanthaceae. *Kew. Bull.* 55: 513–589.
- Stevens, P.F. (2014). [continuously updated]. *Angiosperm Phylogeny Website, version 13*. <http://www.mobot.org/MOBOT/research/APweb/> (accessed on 29 Jun. 2014).
- Xia, C. & Deng, Y.F. (2013). *Phlogacanthus yangtsekiangensis*, a new combination in Chinese Acanthaceae. *Phytotaxa* 104(1): 58–60.

